

# Smarandache’s Cevian Triangle Theorem in The Einstein Relativistic Velocity Model of Hyperbolic Geometry

Cătălin Barbu

“Vasile Alecsandri” College — Bacău, str. Iosif Cocea, nr. 12, sc. A, apt. 13, Romania. E-mail: kafka\_mate@yahoo.com

In this note, we present a proof of Smarandache’s cevian triangle hyperbolic theorem in the Einstein relativistic velocity model of hyperbolic geometry.

## 1 Introduction

Hyperbolic geometry appeared in the first half of the 19<sup>th</sup> century as an attempt to understand Euclid’s axiomatic basis for geometry. It is also known as a type of non-Euclidean geometry, being in many respects similar to Euclidean geometry. Hyperbolic geometry includes such concepts as: distance, angle and both of them have many theorems in common. There are known many main models for hyperbolic geometry, such as: Poincaré disc model, Poincaré half-plane, Klein model, Einstein relativistic velocity model, etc. The hyperbolic geometry is a non-Euclidian geometry. Here, in this study, we present a proof of Smarandache’s cevian triangle hyperbolic theorem in the Einstein relativistic velocity model of hyperbolic geometry. Smarandache’s cevian triangle theorem states that if  $A_1B_1C_1$  is the cevian triangle of point  $P$  with respect to the triangle  $ABC$ , then  $\frac{PA}{PA_1} \cdot \frac{PB}{PB_1} \cdot \frac{PC}{PC_1} = \frac{AB \cdot BC \cdot CA}{A_1B \cdot B_1C \cdot C_1A}$  [1].

Let  $D$  denote the complex unit disc in complex  $z$  - plane, i.e.

$$D = \{z \in \mathbb{C} : |z| < 1\}.$$

The most general Möbius transformation of  $D$  is

$$z \rightarrow e^{i\theta} \frac{z_0 + z}{1 + \bar{z}_0 z} = e^{i\theta} (z_0 \oplus z),$$

which induces the Möbius addition  $\oplus$  in  $D$ , allowing the Möbius transformation of the disc to be viewed as a Möbius left gyrotranslation

$$z \rightarrow z_0 \oplus z = \frac{z_0 + z}{1 + \bar{z}_0 z}$$

followed by a rotation. Here  $\theta \in \mathbb{R}$  is a real number,  $z, z_0 \in D$ , and  $\bar{z}_0$  is the complex conjugate of  $z_0$ . Let  $Aut(D, \oplus)$  be the automorphism group of the grupoid  $(D, \oplus)$ . If we define

$$gyr : D \times D \rightarrow Aut(D, \oplus), \quad gyr[a, b] = \frac{a \oplus b}{b \oplus a} = \frac{1 + \bar{a}b}{1 + \bar{a}b},$$

then is true gyrocommutative law

$$a \oplus b = gyr[a, b](b \oplus a).$$

A gyrovector space  $(G, \oplus, \otimes)$  is a gyrocommutative gyrogroup  $(G, \oplus)$  that obeys the following axioms:

- (1)  $gyr[\mathbf{u}, \mathbf{v}]\mathbf{a} \cdot gyr[\mathbf{u}, \mathbf{v}]\mathbf{b} = \mathbf{a} \cdot \mathbf{b}$  for all points  $\mathbf{a}, \mathbf{b}, \mathbf{u}, \mathbf{v} \in G$ ;

- (2)  $G$  admits a scalar multiplication,  $\otimes$ , possessing the following properties. For all real numbers  $r, r_1, r_2 \in \mathbb{R}$  and all points  $\mathbf{a} \in G$ :

G1  $1 \otimes \mathbf{a} = \mathbf{a}$ ,

G2  $(r_1 + r_2) \otimes \mathbf{a} = r_1 \otimes \mathbf{a} \oplus r_2 \otimes \mathbf{a}$ ,

G3  $(r_1 r_2) \otimes \mathbf{a} = r_1 \otimes (r_2 \otimes \mathbf{a})$ ,

G4  $\frac{|r| \otimes \mathbf{a}}{\|r \otimes \mathbf{a}\|} = \frac{\mathbf{a}}{\|\mathbf{a}\|}$ ,

G5  $gyr[\mathbf{u}, \mathbf{v}](r \otimes \mathbf{a}) = r \otimes gyr[\mathbf{u}, \mathbf{v}]\mathbf{a}$ ,

G6  $gyr[r_1 \otimes \mathbf{v}, r_1 \otimes \mathbf{v}] = I$ ;

- (3) Real vector space structure  $(\|G\|, \oplus, \otimes)$  for the set  $\|G\|$  of onedimensional “vectors”

$$\|G\| = \{\pm \|\mathbf{a}\| : \mathbf{a} \in G\} \subset \mathbb{R}$$

with vector addition  $\oplus$  and scalar multiplication  $\otimes$ , such that for all  $r \in \mathbb{R}$  and  $\mathbf{a}, \mathbf{b} \in G$ :

G7  $\|r \otimes \mathbf{a}\| = |r| \|\mathbf{a}\|$ ,

G8  $\|\mathbf{a} \oplus \mathbf{b}\| \leq \|\mathbf{a}\| \oplus \|\mathbf{b}\|$ .


**Theorem 1 The Hyperbolic Theorem of Ceva in Einstein Gyrovector Space** *Let  $\mathbf{a}_1, \mathbf{a}_2$ , and  $\mathbf{a}_3$  be three non-gyrocollinear points in an Einstein gyrovector space  $(V_s, \oplus, \otimes)$ . Furthermore, let  $\mathbf{a}_{123}$  be a point in their gyroplane, which is off the gyrolines  $\mathbf{a}_1\mathbf{a}_2, \mathbf{a}_2\mathbf{a}_3$ , and  $\mathbf{a}_3\mathbf{a}_1$ . If  $\mathbf{a}_1\mathbf{a}_{123}$  meets  $\mathbf{a}_2\mathbf{a}_3$  at  $\mathbf{a}_{23}$ , etc., then*

$$\frac{\gamma_{\ominus \mathbf{a}_1 \oplus \mathbf{a}_{12}} \|\ominus \mathbf{a}_1 \oplus \mathbf{a}_{12}\|}{\gamma_{\ominus \mathbf{a}_2 \oplus \mathbf{a}_{12}} \|\ominus \mathbf{a}_2 \oplus \mathbf{a}_{12}\|} \cdot \frac{\gamma_{\ominus \mathbf{a}_2 \oplus \mathbf{a}_{23}} \|\ominus \mathbf{a}_2 \oplus \mathbf{a}_{23}\|}{\gamma_{\ominus \mathbf{a}_3 \oplus \mathbf{a}_{23}} \|\ominus \mathbf{a}_3 \oplus \mathbf{a}_{23}\|} \times \frac{\gamma_{\ominus \mathbf{a}_3 \oplus \mathbf{a}_{13}} \|\ominus \mathbf{a}_3 \oplus \mathbf{a}_{13}\|}{\gamma_{\ominus \mathbf{a}_1 \oplus \mathbf{a}_{13}} \|\ominus \mathbf{a}_1 \oplus \mathbf{a}_{13}\|} = 1,$$

(here  $\gamma_v = \frac{1}{\sqrt{1 - \frac{\|v\|^2}{s^2}}}$  is the gamma factor). (See [2, pp. 461].)

**Theorem 2 The Hyperbolic Theorem of Menelaus in Einstein Gyrovector Space** *Let  $\mathbf{a}_1, \mathbf{a}_2$ , and  $\mathbf{a}_3$  be three non-gyrocollinear points in an Einstein gyrovector space  $(V_s, \oplus, \otimes)$ . If a gyroline meets the sides of gyrotriangle  $\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3$  at points  $\mathbf{a}_{12}, \mathbf{a}_{13}, \mathbf{a}_{23}$ , then*

$$\frac{\gamma_{\ominus \mathbf{a}_1 \oplus \mathbf{a}_{12}} \|\ominus \mathbf{a}_1 \oplus \mathbf{a}_{12}\|}{\gamma_{\ominus \mathbf{a}_2 \oplus \mathbf{a}_{12}} \|\ominus \mathbf{a}_2 \oplus \mathbf{a}_{12}\|} \cdot \frac{\gamma_{\ominus \mathbf{a}_2 \oplus \mathbf{a}_{23}} \|\ominus \mathbf{a}_2 \oplus \mathbf{a}_{23}\|}{\gamma_{\ominus \mathbf{a}_3 \oplus \mathbf{a}_{23}} \|\ominus \mathbf{a}_3 \oplus \mathbf{a}_{23}\|} \times \frac{\gamma_{\ominus \mathbf{a}_3 \oplus \mathbf{a}_{13}} \|\ominus \mathbf{a}_3 \oplus \mathbf{a}_{13}\|}{\gamma_{\ominus \mathbf{a}_1 \oplus \mathbf{a}_{13}} \|\ominus \mathbf{a}_1 \oplus \mathbf{a}_{13}\|} = 1.$$



(See [2, pp. 463].) For further details we refer to A. Ungar’s recent book [2].

**2 Main result**

In this section, we present a proof of Smarandache’s cevian triangle hyperbolic theorem in the Einstein relativistic velocity model of hyperbolic geometry.

**Theorem 3** *If  $A_1B_1C_1$  is the cevian gyrotriangle of gyropoint  $P$  with respect to the gyrotriangle  $ABC$ , then*

$$\frac{\gamma_{|PA_1||PA_1|} \cdot \gamma_{|PB_1||PB_1|} \cdot \gamma_{|PC_1||PC_1|}}{\gamma_{|PA_1||PA_1|} \cdot \gamma_{|PB_1||PB_1|} \cdot \gamma_{|PC_1||PC_1|}} = \frac{\gamma_{|AB_1||AB_1|} \cdot \gamma_{|BC_1||BC_1|} \cdot \gamma_{|CA_1||CA_1|}}{\gamma_{|AB_1||AB_1|} \cdot \gamma_{|BC_1||BC_1|} \cdot \gamma_{|CA_1||CA_1|}}$$

**Proof** *If we use a theorem 2 in the gyrotriangle  $ABC$  (see Figure), we have*

$$\gamma_{|AC_1||AC_1|} \cdot \gamma_{|BA_1||BA_1|} \cdot \gamma_{|CB_1||CB_1|} = \gamma_{|AB_1||AB_1|} \cdot \gamma_{|BC_1||BC_1|} \cdot \gamma_{|CA_1||CA_1|} \quad (1)$$

*If we use a theorem 1 in the gyrotriangle  $AA_1B$ , cut by the gyroline  $CC_1$ , we get*

$$\gamma_{|AC_1||AC_1|} \cdot \gamma_{|BC_1||BC_1|} \cdot \gamma_{|A_1P||A_1P|} = \gamma_{|AP||AP|} \cdot \gamma_{|A_1C_1||A_1C_1|} \cdot \gamma_{|BC_1||BC_1|} \quad (2)$$

*If we use a theorem 1 in the gyrotriangle  $BB_1C$ , cut by the gyroline  $AA_1$ , we get*

$$\gamma_{|BA_1||BA_1|} \cdot \gamma_{|CA_1||CA_1|} \cdot \gamma_{|B_1P||B_1P|} = \gamma_{|BP||BP|} \cdot \gamma_{|B_1A_1||B_1A_1|} \cdot \gamma_{|CA_1||CA_1|} \quad (3)$$

*If we use a theorem 1 in the gyrotriangle  $CC_1A$ , cut by the gyroline  $BB_1$ , we get*

$$\gamma_{|CB_1||CB_1|} \cdot \gamma_{|AB_1||AB_1|} \cdot \gamma_{|C_1P||C_1P|} = \gamma_{|CP||CP|} \cdot \gamma_{|C_1B_1||C_1B_1|} \cdot \gamma_{|AB_1||AB_1|} \quad (4)$$

*We divide each relation (2), (3), and (4) by relation (1), and we obtain*

$$\frac{\gamma_{|PA_1||PA_1|}}{\gamma_{|PA_1||PA_1|}} = \frac{\gamma_{|BC_1||BC_1|}}{\gamma_{|BA_1||BA_1|}} \cdot \frac{\gamma_{|B_1A_1||B_1A_1|}}{\gamma_{|B_1C_1||B_1C_1|}}, \quad (5)$$

$$\frac{\gamma_{|PB_1||PB_1|}}{\gamma_{|PB_1||PB_1|}} = \frac{\gamma_{|CA_1||CA_1|}}{\gamma_{|CB_1||CB_1|}} \cdot \frac{\gamma_{|C_1B_1||C_1B_1|}}{\gamma_{|C_1A_1||C_1A_1|}}, \quad (6)$$

$$\frac{\gamma_{|PC_1||PC_1|}}{\gamma_{|PC_1||PC_1|}} = \frac{\gamma_{|AB_1||AB_1|}}{\gamma_{|AC_1||AC_1|}} \cdot \frac{\gamma_{|A_1C_1||A_1C_1|}}{\gamma_{|A_1B_1||A_1B_1|}} \quad (7)$$

*Multiplying (5) by (6) and by (7), we have*

$$\begin{aligned} & \frac{\gamma_{|PA_1||PA_1|}}{\gamma_{|PA_1||PA_1|}} \cdot \frac{\gamma_{|PB_1||PB_1|}}{\gamma_{|PB_1||PB_1|}} \cdot \frac{\gamma_{|PC_1||PC_1|}}{\gamma_{|PC_1||PC_1|}} = \\ & \frac{\gamma_{|AB_1||AB_1|} \cdot \gamma_{|BC_1||BC_1|} \cdot \gamma_{|CA_1||CA_1|}}{\gamma_{|AB_1||AB_1|} \cdot \gamma_{|BC_1||BC_1|} \cdot \gamma_{|CA_1||CA_1|}} \cdot \frac{\gamma_{|B_1A_1||B_1A_1|} \cdot \gamma_{|C_1B_1||C_1B_1|} \cdot \gamma_{|A_1C_1||A_1C_1|}}{\gamma_{|A_1B_1||A_1B_1|} \cdot \gamma_{|B_1C_1||B_1C_1|} \cdot \gamma_{|C_1A_1||C_1A_1|}} \end{aligned} \quad (8)$$

*From the relation (1) we have*

$$\frac{\gamma_{|B_1A_1||B_1A_1|} \cdot \gamma_{|C_1B_1||C_1B_1|} \cdot \gamma_{|A_1C_1||A_1C_1|}}{\gamma_{|A_1B_1||A_1B_1|} \cdot \gamma_{|B_1C_1||B_1C_1|} \cdot \gamma_{|C_1A_1||C_1A_1|}} = 1, \quad (9)$$

so

$$\frac{\gamma_{|PA_1||PA_1|} \cdot \gamma_{|PB_1||PB_1|} \cdot \gamma_{|PC_1||PC_1|}}{\gamma_{|PA_1||PA_1|} \cdot \gamma_{|PB_1||PB_1|} \cdot \gamma_{|PC_1||PC_1|}} = \frac{\gamma_{|AB_1||AB_1|} \cdot \gamma_{|BC_1||BC_1|} \cdot \gamma_{|CA_1||CA_1|}}{\gamma_{|AB_1||AB_1|} \cdot \gamma_{|BC_1||BC_1|} \cdot \gamma_{|CA_1||CA_1|}}$$

Submitted on March 05, 2010 / Accepted on March 26, 2010

**References**

1. Smarandache F. Eight solved and eight open problems in elementary geometry. arXiv: 1003.2153.
2. Ungar A.A. Analytic hyperbolic geometry and Albert Einstein’s Special Theory of Relativity. World Scientific Publishing Co., Hackensack (NJ), 2008.