

# Type III Spacetime with Closed Timelike Curves

Ahmed Faizuddin

Hindustani Kendriya Vidyalaya, Dinseh Ojha Road, Bhangagarh, Guwahati-05, Assam, India  
E-mail: faizuddinahmed15@gmail.com

We present a symmetric spacetime, admitting closed timelike curves (CTCs) which appear after a certain instant of time, *i.e.*, a time-machine spacetime. These closed timelike curves evolve from an initial spacelike hypersurface on the planes  $z = \text{constant}$  in a causally well-behaved manner. The spacetime discussed here is free from curvature singularities and a 4D generalization of the Misner space in curved spacetime. The matter field is of pure radiation with cosmological constant.

## 1 Introduction

One of the most intriguing aspects of Einstein's theory of gravitation is that solutions of field equations admit closed timelike curves (CTC). Presence of CTC in a spacetime leads to time-travel which violates the causality condition. The first one being Gödel's spacetime [1] which admits closed timelike curves (CTC) everywhere and an eternal time-machine spacetime. There are a considerable number of spacetimes in literature that admitting closed timelike curves have been constructed. A small sample would be [1–21]. One way of classifying such causality violating spacetimes would be to categorize the metrics as either eternal time-machine in which CTC always exist (in this class would be [1, 2]), or as time-machine spacetimes in which CTC appear after a certain instant of time. In the latter category would be the ones discussed in [18–20]. Many of the models, however, suffer from one or more severe drawbacks. For instance, in some of these solutions, for example [13, 14, 20], the weak energy condition (WEC) is violated indicating unrealistic matter-energy content and some other solutions have singularities.

Among the time-machine spacetimes, we mention two: the first being Ori's compact core [17] which is represented by a vacuum metric locally isometric to  $pp$  waves and second, which is more relevant to the present work, the Misner space [22] in 2D. This is essentially a two dimensional metric (hence flat) with peculiar identifications. The Misner space is interesting in the context of CTC as it is a prime example of a spacetime where CTC evolve from causally well-behaved initial conditions.

The metric for the Misner space [22]

$$ds_{\text{Misn}}^2 = -2 dt dx - t dx^2 \quad (1)$$

where  $-\infty < t < \infty$  but the co-ordinate  $x$  is periodic. The metric (1) is regular everywhere as  $\det g = -1$  including at  $t = 0$ . The curves  $t = t_0$ , where  $t_0$  is a constant, are closed since  $x$  is periodic. The curves  $t < 0$  are spacelike, but  $t > 0$  are timelike and the null curves  $t = 0$  form the chronology horizon. The second type of curves, namely,  $t = t_0 > 0$  are closed timelike curves (CTC). This metric has been the subject of intense study and quite recently, Levanony and Ori [23], have studied the motion of extended bodies in the

2D Misner space and its flat 4D generalizations. A non-flat 4D spacetime, satisfying all the energy conditions, but with causality violating properties of the Misner space, primarily that CTC evolve smoothly from an initially causally well-behaved stage, would be physically more acceptable as a time-machine spacetime.

In this paper, we shall attempt to show that causality violating curves appear in non-vacuum spacetime with comparatively simple structure. In section 2, we analyze the spacetime; in section 3, the matter distribution and energy condition; in section 4, the spacetime is classified and its kinematical properties discussed; and concluding in section 5.

## 2 Analysis of the spacetime

Consider the following metric

$$ds^2 = 4r^2 dr^2 + e^{2\alpha r^2} (dz^2 - t d\phi^2 - 2 dt d\phi) + 4\beta z r e^{-\alpha r^2} dr d\phi \quad (2)$$

where  $\phi$  coordinate is assumed periodic  $0 \leq \phi \leq \phi_0$ , where  $\alpha$  is an integer and  $\beta > 0$  is a real number. We have used co-ordinates  $x^1 = r$ ,  $x^2 = \phi$ ,  $x^3 = z$  and  $x^4 = t$ . The ranges of the other co-ordinates are  $t, z \in (-\infty, \infty)$  and  $0 \leq r < \infty$ . The metric has signature  $(+, +, +, -)$  and the determinant of the corresponding metric tensor  $g_{\mu\nu}$ ,  $\det g = -4r^2 e^{6\alpha r^2}$ . The non-zero components of the Einstein tensor are

$$G_{\mu}^{\mu} = 3\alpha^2, \quad G_{\phi}^t = -\frac{1}{2} e^{-6\alpha r^2} \beta^2. \quad (3)$$

Consider an azimuthal curve  $\gamma$  defined by  $r = r_0$ ,  $z = z_0$  and  $t = t_0$ , where  $r_0, z_0, t_0$  are constants, then we have from the metric (2)

$$ds^2 = -t e^{2\alpha r^2} d\phi^2. \quad (4)$$

These curves are null for  $t = 0$ , spacelike throughout for  $t = t_0 < 0$ , but become timelike for  $t = t_0 > 0$ , which indicates the presence of closed timelike curves (CTC). Hence CTC form at a definite instant of time satisfy  $t = t_0 > 0$ .

It is crucial to have analysis that the above CTC evolve from a spacelike  $t = \text{constant}$  hypersurface (and thus  $t$  is a

time coordinate) [17]. This can be ascertained by calculating the norm of the vector  $\nabla_\mu t$  (or by determining the sign of the component  $g^{tt}$  in the inverse metric tensor  $g^{\mu\nu}$  [17]). We find from (2) that

$$g^{tt} = t e^{-2\alpha r^2} + \beta^2 z^2 e^{-6\alpha r^2}. \tag{5}$$

A hypersurface  $t = \text{constant}$  is spacelike provided  $g^{tt} < 0$  for  $t = t_0 < 0$ , but becomes timelike provided  $g^{tt} > 0$  for  $t = t_0 > 0$ . Here we choose the  $z$ -planes defined by  $z = z_0$ , ( $z_0$ , a constant equal to zero) such that the above condition is satisfied. Thus the spacelike  $t = \text{constant} < 0$  hypersurface can be chosen as initial conditions over which the initial may be specified. There is a Cauchy horizon for  $t = t_0 = 0$  called Chronology horizon which separates the causal and non-causal parts of the spacetime. Hence the spacetime evolves from a partial Cauchy hypersurface (initial spacelike hypersurface) in a causally well-behaved manner, up to a moment, *i.e.*, a null hypersurface  $t = 0$  and CTC form at a definite instant of time on  $z = \text{constant}$  plane.

Consider the Killing vector  $\eta = \partial_\phi$  for metric (2) which has the normal form

$$\eta^\mu = (0, 1, 0, 0). \tag{6}$$

Its co-vector is

$$\eta_\mu = (2\beta z r e^{-\alpha r^2}, -t e^{2\alpha r^2}, 0, -e^{2\alpha r^2}). \tag{7}$$

The (6) satisfies the Killing equation  $\eta_{\mu;\nu} + \eta_{\nu;\mu} = 0$ . For cyclicly symmetric metric, the norm  $\eta_\mu \eta^\mu$  of the Killing vector is spacelike, closed orbits [24–28]. We note that

$$r^\mu \eta_\mu = -t e^{2\alpha r^2} \tag{8}$$

which is spacelike for  $t < 0$ , closed orbits ( $\phi$  co-ordinate being periodic).

An important note is that the Riemann tensor  $R_{\mu\nu\rho\sigma}$  can be expressed in terms of the metric tensor  $g_{\mu\nu}$  as

$$R_{\mu\nu\rho\sigma} = k (g_{\mu\rho} g_{\nu\sigma} - g_{\mu\sigma} g_{\nu\rho}) \tag{9}$$

where  $k = -\alpha^2$  for the spacetime (2).

Another important note is that if we take  $\beta = 0$ , then the spacetime represented by (2) is maximally symmetric vacuum spacetime and locally isometric anti-de Sitter space in four-dimension. One can easily show by a number of transformations the standard form of locally isometric  $AdS_4$  metric [29]

$$ds^2 = \frac{3}{(-\Lambda) x^2} (-dt^2 + dx^2 + d\phi^2 + dz^2) \tag{10}$$

where one of the co-ordinate  $\phi$  being periodic.

### 3 Matter distribution of the spacetime and energy condition

Einstein’s field equations taking into account the cosmological constant

$$G^{\mu\nu} + \Lambda g^{\mu\nu} = T^{\mu\nu}, \quad \mu, \nu = 1, 2, 3, 4. \tag{11}$$

Consider the energy-momentum tensor of pure radiation field [30]

$$T^{\mu\nu} = \rho n^\mu n^\nu \tag{12}$$

where  $n^\mu$  is the null vector defined by

$$n^\mu = (0, 0, 0, 1). \tag{13}$$

The non-zero component of the energy-momentum tensor

$$T^t_\phi = -\rho e^{2\alpha r^2}. \tag{14}$$

Equating field equations (11) using (3) and (14), we get

$$\begin{aligned} \Lambda &= -3\alpha^2, \\ \rho &= \frac{1}{2} \beta^2 e^{-8\alpha r^2}, \quad 0 \leq r < \infty. \end{aligned} \tag{15}$$

The energy-density of pure radiation or null dust decreases exponentially with  $r$  and vanish at  $r \rightarrow \pm\infty$ . The matter field pure radiation satisfy the energy condition and the energy density  $\rho$  is always positive.

### 4 Classification and kinematical properties of the spacetime

For classification of the spacetime (2), we can construct the following set of null tetrads ( $k, l, m, \bar{m}$ ) as

$$k_\mu = (0, 1, 0, 0), \tag{16}$$

$$l_\mu = \left(-2\beta z r e^{-\alpha r^2}, \frac{t}{2} e^{2\alpha r^2}, 0, e^{2\alpha r^2}\right), \tag{17}$$

$$m_\mu = \frac{1}{\sqrt{2}} (2r, 0, i e^{\alpha r}, 0), \tag{18}$$

$$\bar{m}_\mu = \frac{1}{\sqrt{2}} (2r, 0, -i e^{\alpha r}, 0), \tag{19}$$

where  $i = \sqrt{-1}$ . The set of null tetrads above are such that the metric tensor for the line element (2) can be expressed as

$$g_{\mu\nu} = -k_\mu l_\nu - l_\mu k_\nu + m_\mu \bar{m}_\nu + \bar{m}_\mu m_\nu. \tag{20}$$

The vectors (16)–(19) are null vectors and are orthogonal except for  $k_\mu l^\mu = -1$  and  $m_\mu \bar{m}^\mu = 1$ . Using this null tetrad above, we have calculated the five Weyl scalars

$$\begin{aligned} \Psi_3 &= -\frac{i\alpha\beta e^{-2\alpha r^2}}{2\sqrt{2}}, \\ \Psi_4 &= -\frac{1}{4} \beta e^{-2\alpha r^2} (i + 2\alpha z e^{\alpha r^2}) \end{aligned} \tag{21}$$

are non-vanishing, while  $\Psi_0 = \Psi_1 = \Psi_2 = 0$ . The spacetime represented by (2) is of type III in the Petrov classification scheme. Note that the non-zero Weyl scalars  $\Psi_3$  and  $\Psi_4$  are finite at  $r \rightarrow 0$  and vanish as  $r \rightarrow \pm\infty$  indicating asymptotic flatness of the spacetime (2). The metric (2) is free

from curvature singularities. The curvature invariant known as Kretschmann scalar is given by

$$R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} = 24 \alpha^4 \quad (22)$$

and the curvature scalar

$$R = -12 \alpha^2 \quad (23)$$

are constant being non-zero.

Using the null tetrad (16) we have calculated the *Optical* scalars [30] the *expansion*, the *twist* and the *shear* and they are

$$\begin{aligned} \Theta &= \frac{1}{2} k^\mu_{;\mu} = 0, \\ \omega^2 &= \frac{1}{2} k_{[\mu;\nu]} k^{\mu;\nu} = 0, \\ \sigma \bar{\sigma} &= \frac{1}{2} k_{(\mu;\nu)} k^{\mu;\nu} - \Theta^2 = 0 \end{aligned} \quad (24)$$

and the null vector (16) satisfy the geodesics equation

$$k_{\mu;\nu} k^\nu = 0. \quad (25)$$

Thus the spacetime represented by (2) is non-diverging, has shear-free null geodesics congruence. One can easily show that for constant  $r$  and  $z$ , the metric (2) reduces to conformal Misner space in 2D

$$ds_{confo}^2 = \Omega ds_{Mishn}^2 \quad (26)$$

where  $\Omega = e^{2\alpha r^2}$  is a constant.

## 5 Conclusion

Our primary motivation in this paper is to write down a metric for a spacetime that incorporates the Misner space and its causality violating properties and to classify it. The solution presented here is non-vacuum, cyclicly symmetric metric (2) and serves as a model of time-machine spacetime in the sense that CTC appear at a definite instant of time on the  $z$ -plane. Most of the CTC spacetimes violate one or more energy conditions or unrealistic matter source and are unphysical. The model discussed here is free from all these problems and matter distribution is of pure radiation field with negative cosmological constant satisfying the energy condition.

Submitted on July 25, 2016 / Accepted on August 9, 2016

## References

- Godel K. An example of a new type of cosmological solutions of Einsteins field equations of gravitation. *Rev. Mod. Phys.*, 1949, v.21 (3), 447–450.
- van Stockum W.J. The gravitational field of a distribution of particles rotating about an axis of symmetry. *Proc. R. Soc. Edin.*, 1937, v.57, 135–154.
- Tipler F.J. Rotating Cylinders and the Possibility of Global Causality Violation. *Phys. Rev. D*, 1974, v.9 (8), 2203–2206.
- Gott J.R. Closed timelike curves produced by pairs of moving cosmic strings: Exact solutions. *Phys. Rev. Lett.*, 1991, v.66 (9), 1126–1129.
- Bonnor W.B. An exact, asymptotically flat, vacuum solution of Einstein's equations with closed timelike curves. *Class. Quantum Grav.*, 2002, v.19, 5951–5957.
- Bonnor W.B. Closed timelike curves in general relativity. *Int. J. Mod. Phys. D*, 2003, v.12, 1705–1708.
- Bonnor W.B., Steadman B.R. Exact solutions of the Einstein Maxwell equations with closed timelike curves. *Gen. Rel. Grav.*, 2005, v.37 (11), 1833–1844.
- Kerr R.P. Gravitational Field of a Spinning Mass as an Example of Algebraically Special Metrics. *Phys. Rev. Lett.*, 1963, v.11 (5), 237–238.
- Carter B. Global structure of the Kerr family of gravitational fields. *Phys. Rev. D*, 1968, v.174, 1559–1571.
- Gürses M., Karasu A., Sarioğlu Ö. Godel-type metrics in various dimensions. *Class. Quantum Grav.*, 2005, v.22 (9), 1527–1543.
- Alcubierre M. The warp drive: hyper-fast travel within general relativity. *Class. Quantum Grav.*, 1994, v.11, L73–L77.
- Lobo F.S.N., Crawford P. Time, closed timelike curves and causality. *The Nature of Time : Geometry, Physics and Perception. NATO Sci. Ser. II*, 2003, v.95, 289–296.
- Morris M.S., Thorne K.S., Yurtsever U. Time Machines and the Weak Energy Conditions. *Phys. Rev. Lett.*, 1988, v.61 (13), 1446–1449.
- Morris M.S., Thorne K.S. Wormholes in space-time and their use for interstellar travel: A tool for teaching general relativity. *Amer. J. Phys.*, 1988, v.56, 395–412.
- Krasnikov S.V. Hyper-fast interstellar travel in general relativity. *Phys. Rev. D*, 1998, v.57 (8), 4760–4766.
- Everett A.E., Roman T.A. A superluminal subway: the Krasnikov tube. *Phys. Rev. D*, 1997, v.56, 2100–2108. arXiv: gr-qc/9702049.
- Ori A. A new time-machine model with compact vacuum core. *Phys. Rev. Lett.*, 2005, v.95, 021101–021107.
- Ori A. Formation of closed timelike curves in a composite vacuum/dust asymptotically-flat spacetime. *Phys. Rev. D*, 2007, v.76, 044002–044015.
- Ori A. Improved time-machine model. *Phys. Rev. D*, 1996, v.54 (8), 4858–4861.
- Ori A. Causality violation and the weak energy condition. *Phys. Rev. D*, 1994, v.49 (8), 3990–3997.
- Sarma D., Patgiri M., Ahmed F.U. Pure radiation metric with stable closed timelike curves. *Gen. Rel. Grav.*, 2014, v.46, 1633–1641.
- Misner C.W. Taub-NUT Space as a Counterexample to Almost Anything. In Ehlers J., ed. *Relativity Theory and Astrophysics 1. Relativity and Cosmology*, Am. Math. Soc., Providence, RI, 1967, 160–169.
- Levanony D., Ori A. Extended time-travelling object in Misner space. *Phys. Rev. D*, 2011, v.83, 044043. arXiv: gr-qc/1102.0907.
- Mars M., Senovilla J.M.M. Axial symmetry and conformal Killing vectors. *Class. Quantum Grav.*, 1993, v.10, 1633–1647.
- Carter B. The commutation property of a stationary, axisymmetric system. *Commun. Math. Phys.*, 1970, v.17, 233–238.
- Branes A. Some restrictions on the symmetry groups of axially symmetric spacetimes. *Class. Quantum Grav.*, 2001, v.18, 5511–5520.
- Bičák J., Schmidt B.D. Isometries compatible with gravitational radiation. *J. Math. Phys.*, 1984, v.25 (3), 600–606.
- Wald R.M. *General Relativity*. Cambridge University Press, Cambridge, 1984.
- Zofka M., Bičák J. Cylindrical spacetimes with  $\Lambda \neq 0$  and their sources. *Class. Quantum Grav.*, 2008, v.25 (1), 015011. arXiv: gr-qc/0712.2144
- Stephani H., Kramer D., MacCallum M., Hoenselaers C., Herlt E. *Exact Solutions to Einstein's Field Equations*, Cambridge Univ. Press, 2003.